


US Serial No. 10/597404
Page 2 of 25

RECEIVED
CENTRAL FAX CENTER
APR 28 2009

In the Claims:

1.(Currently amended) A process for preparing compositions which provide upon activation an olfactory compound comprising:

A) incorporating into the composition a compound of formula (I) as precursor for olfactory compounds compound



wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;

Y is $-CR^5R^6R^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($R^5 + R^6 + R^7$) is not greater than 18; or

Y is $-CR^5R^6R^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ($R^5 + R^6 + R^7$) is not greater than 18; or

US Serial No. 10/597404
Page 3 of 25

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18; or

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18;


R^2 and R^3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue, $-\text{NH}_2$, $-\text{NO}_2$, $-\text{NHCO}_2\text{CH}_3$, $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$, $-\text{N}(\text{hydroxyalkyl})_2$, $-\text{NHC}(\text{O})-(\text{C}_1\text{-C}_8 \text{ alkyl})$ or $-\text{NHC}(\text{O})-(\text{C}_3\text{-C}_8 \text{ aryl})$; or

R^2 and R^3 are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

R^4 in 2- or 3-position is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_3 - C_6 cycloalkyl, or $-\text{CN}$; and

- a) if n is zero, R is a C_1 - C_{24} hydrocarbon residue, or C_1 - C_{24} hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or
- b) if n is 1, R is a C_1 - C_{25} hydrocarbon residue, a C_1 - C_{25} hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O), or C_1 - C_{25} hydrocarbon residue substituted by an ionic substituent of the formula $\text{N}(\text{R}^{20})_3^+$, in which R^{20} is the residue of an alkyl group with 1 to 18 carbon atoms; or
- R is a monovalent residue of the formula (i)

US Serial No. 10/597404
Page 4 of 25



wherein

X is $-\text{CR}^{14}\text{R}^{15}\text{R}^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($\text{R}^{14} + \text{R}^{15} + \text{R}^{16}$) is not greater than 18; or

X is $-\text{CR}^{14}\text{R}^{15}\text{R}^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ($\text{R}^{14} + \text{R}^{15} + \text{R}^{16}$) is not greater than 18; or

X is $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$) is not greater than 18; or

X is $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$) is not greater than 18;

R^{12} and R^{13} are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue,

US Serial No. 10/597404
Page 5 of 25

-NO₂, -NH₂, -NHCO₂CH₃, -N(C₁-C₆ alkyl)₂, -N(hydroxyalkyl)₂,
-NHC(O)-(C₁-C₈ alkyl) or -NHC(O)-(C₃-C₈ aryl); or

R¹² and R¹³ are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix),
and form together with the carbon atoms to which they are attached a
dioxolane ring or a dioxane ring;

R¹¹ in ii- or iii-position is hydrogen, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₃-C₆
cycloalkyl, or -CN; and

B) cleaving the compound of formula (I) by hydrolysis followed by
photoisomerization.

2. (canceled)

3. (canceled)

4.(canceled)

5.(canceled)

6.(previously presented) A process according to claim 1 for preparing compositions
which provide upon activation an olfactory compound comprising incorporating into
the composition a compound of formula (I) wherein R⁵, R⁶ and R⁷ are independently
hydrogen or a C₁-C₁₀ hydrocarbon residue.

7.(previously presented) A process according to claim 1 wherein the sum of carbon
atoms R⁵ + R⁶ + R⁷ is between 6 and 15.

US Serial No. 10/597404
Page 6 of 25

8.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein at least one residue R^5 , R^6 and R^7 is not hydrogen.

9.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{10} hydrocarbon residue.

10.(previously presented) A process according to claim 1 wherein the sum of carbon atoms $R^8 + R^9 + R^{10}$ is between 6 and 15.

11.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein at least one residue R^8 , R^9 and R^{10} is not hydrogen.

12.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{10} hydrocarbon residue.

13.(previously presented) A process according to claim 1 wherein the sum of carbon atoms $R^{14} + R^{15} + R^{16}$ is between 6 and 15.

14.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein at least one residue R^{14} , R^{15} and R^{16} is not hydrogen.

US Serial No. 10/597404
Page 7 of 25

15.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{10} hydrocarbon residue.

16.(previously presented) A process according to claim 1 wherein the sum of carbon atoms $R^{17} + R^{18} + R^{19}$ is between 6 and 15.

17.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein at least one residue R^{17} , R^{18} and R^{19} is not hydrogen.

18.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein n is 1.

19.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein n is 1 and Y a residue selected from the group consisting of a fragrant alcohol $HO-CR^5R^6R^7$, of the enol form of a fragrant aldehyde of the formula $O=(CH)-CHR^9R^{10}$, and of the enol form of a fragrant ketone of the formula $O=(CR^8)-CHR^9R^{10}$.

20.(previously presented) A process according to claim 19 wherein the fragrant alcohol having a molecular weight between 46 and 400.

21.(previously presented) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into

US Serial No. 10/597404
Page 8 of 25

the composition a compound of formula (I) wherein n is 1 and Y a residue selected from the group consisting of a fragrant alcohol $\text{HO}-\text{CR}^5\text{R}^6\text{R}^7$, of the enol form of a fragrant aldehyde of the formula $\text{O}=(\text{CH})-\text{CHR}^9\text{R}^{10}$, and of the enol form of a fragrant ketone of the formula $\text{O}=(\text{CR}^8)-\text{CHR}^9\text{R}^{10}$, and R is selected from the group consisting of methyl, ethyl, propyl, butyl, pentyl, 2-ethylhexyl, cyclopentyl, cyclohexyl, and the residue of a fragrant alcohol.

22.(previously presented) A process according to claim 21 wherein the fragrant alcohol having a molecular weight between 46 and 400.

23.(previously presented) A process according to claim 19 wherein the fragrant aldehyde having a molecular weight between 100 and 450.

24.(previously presented) A process according to claim 21 wherein the fragrant aldehyde having a molecular weight between 100 and 450.

25.(previously presented) A process according to claim 19 wherein the fragrant ketone having a molecular weight between 100 and 450.

26.(previously presented) A process according to claim 21 wherein the fragrant ketone having a molecular weight between 100 and 450.

27.(currently amended) A process according to claim 1 for preparing compositions which provide upon activation an olfactory compound comprising incorporating into the composition a compound of formula (I) wherein n is 1 and R is a monovalent ~~amonovalent~~ residue of formula (i).

28.(withdrawn) A consumer product comprising a compound of formula (I) as defined by claim 1.


US Serial No. 10/597404
Page 9 of 25

29.(withdrawn) A consumer product according to claim 28 comprising a compound of formula (I) wherein n is 1 and Y a residue selected from the group consisting of a fragrant alcohol $\text{HO}-\text{CR}^5\text{R}^6\text{R}^7$, of the enol form of a fragrant aldehyde of the formula $\text{O}=(\text{CH})-\text{CHR}^9\text{R}^{10}$, and of the enol form of a fragrant ketone of the formula $\text{O}=(\text{CR}^8)-\text{CHR}^9\text{R}^{10}$.

30.(withdrawn) A consumer product according to claim 28 wherein the consumer product is selected from fine fragrance, industrial care products, institutional care products, home care products, personal care products, industrial cleaning products, institutional cleaning products, home cleaning products, and fabric conditioner.

31.(withdrawn-currently amended) A process of providing an olfactory compound to a substrate comprising the steps:

- a) cleaving a compound of formula (I) by hydrolysis followed by photoisomerization



(I)

wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;

US Serial No. 10/597404
Page 10 of 25

Y is $-\text{CR}^5\text{R}^6\text{R}^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($\text{R}^5 + \text{R}^6 + \text{R}^7$) is not greater than 18; or

Y is $-\text{CR}^5\text{R}^6\text{R}^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ($\text{R}^5 + \text{R}^6 + \text{R}^7$) is not greater than 18; or

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18; or

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18;


R^2 and R^3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue, $-\text{NH}_2$, $-\text{NO}_2$, $-\text{NHCO}_2\text{CH}_3$, $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$, $-\text{N}(\text{hydroxyalkyl})_2$, $-\text{NHC}(\text{O})\text{-(C}_1\text{-C}_8 \text{ alkyl)}$ or $-\text{NHC}(\text{O})\text{-(C}_3\text{-C}_8 \text{ aryl)}$;
or

R^2 and R^3 are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

US Serial No. 10/597404
Page 11 of 25

R^4 in 2- or 3-position is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_3 - C_6 cycloalkyl, or $-CN$; and

- a) if n is zero, R is a C_1 - C_{24} hydrocarbon residue, or C_1 - C_{24} hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or
- b) if n is 1, R is a C_1 - C_{25} hydrocarbon residue, a C_1 - C_{25} hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and $C(O)$, or C_1 - C_{25} hydrocarbon residue substituted by an ionic substituent of the formula $N(R^{20})_3^+$, in which R^{20} is the residue of an alkyl group with 1 to 18 carbon atoms; or R is a monovalent residue of the formula (i)



wherein

X is $-CR^{14}R^{15}R^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($R^{14} + R^{15} + R^{16}$) is not greater than 18; or

X is $-CR^{14}R^{15}R^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and $C(O)$, and the sum of all carbon atoms ($R^{14} + R^{15} + R^{16}$) is not greater than 18; or

US Serial No. 10/597404
Page 12 of 25

X is $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$) is not greater than 18; or

X is $-\text{CR}^{17}=\text{CR}^{18}\text{R}^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^{17} + \text{R}^{18} + \text{R}^{19}$) is not greater than 18;


R^{12} and R^{13} are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue, $-\text{NO}_2$, $-\text{NH}_2$, $-\text{NHCO}_2\text{CH}_3$, $-\text{N}(\text{C}_1\text{-C}_6 \text{ alkyl})_2$, $-\text{N}(\text{hydroxyalkyl})_2$, $-\text{NHC}(\text{O})-(\text{C}_1\text{-C}_8 \text{ alkyl})$ or $-\text{NHC}(\text{O})-(\text{C}_3\text{-C}_8 \text{ aryl})$; or

R^{12} and R^{13} are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

R^{11} in ii- or iii-position is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_3 - C_6 cycloalkyl, or $-\text{CN}$;


resulting in a compound of formula (Ia)

US Serial No. 10/597404
Page 13 of 25




wherein R^2 , R^3 , R^4 and Y have the same meaning as given above;
followed by

- b) cleaving the compound of formula (Ia) of step a under activating conditions in the presence of light resulting in a coumarin (IIa)



wherein R^2 , R^3 and R^4 have the same meaning as given above.

32.(withdrawn) A compound of formula (I)



wherein the acrylic acid ester double bond is of the E configuration;

n is zero or 1;

US Serial No. 10/597404
Page 14 of 25

Y is $-\text{CR}^5\text{R}^6\text{R}^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($\text{R}^5 + \text{R}^6 + \text{R}^7$) is not greater than 18 and at least 6; or

Y is $-\text{CR}^5\text{R}^6\text{R}^7$, wherein R^5 , R^6 and R^7 are independently hydrogen or a C_1 - C_{18} aliphatic residue containing one or more atoms/groups selected from O, N and C(O), the sum of all carbon atoms ($\text{R}^5 + \text{R}^6 + \text{R}^7$) is not greater than 18; and at least one of the residues R^5 , R^6 and R^7 is not hydrogen; or

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18; or

Y is $-\text{CR}^8=\text{CR}^9\text{R}^{10}$, wherein R^8 , R^9 and R^{10} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($\text{R}^8 + \text{R}^9 + \text{R}^{10}$) is not greater than 18;


R^2 and R^3 are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue, $-\text{NH}_2$, $-\text{NO}_2$, $-\text{NHCO}_2\text{CH}_3$, $-\text{N}(\text{C}_1\text{-C}_6\text{ alkyl})_2$, $-\text{N}(\text{hydroxyalkyl})_2$, $-\text{NHC}(\text{O})-(\text{C}_1\text{-C}_8\text{ alkyl})$ or $-\text{NHC}(\text{O})-(\text{C}_3\text{-C}_8\text{ aryl})$; or

R^2 and R^3 are attached at the positions C(6,7), C(7,8), or C(8,9), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

R^4 in 2- or 3-position is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_3 - C_6 cycloalkyl, or $-\text{CN}$; and

US Serial No. 10/597404
Page 15 of 25

- a) if n is zero, R is a C_2 - C_{24} hydrocarbon residue, or C_1 - C_{24} hydrocarbon residue containing one or more heteroatoms selected from N, O and Si; or
- b) if n is 1, R is a C_1 - C_{25} hydrocarbon residue, a C_1 - C_{25} hydrocarbon residue containing one or more atoms/groups selected from N, O, Si, and C(O), or C_1 - C_{25} hydrocarbon residue substituted by an ionic substituent of the formula $N(R^{20})_3^+$, in which R^{20} is the residue of an alkyl group with 1 to 18 carbon atoms; or
- R is a monovalent residue of the formula (i)



wherein

X is $-CR^{14}R^{15}R^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, and the sum of all carbon atoms ($R^{14} + R^{15} + R^{16}$) is not greater than 18; or

X is $-CR^{14}R^{15}R^{16}$, wherein R^{14} , R^{15} and R^{16} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), and the sum of all carbon atoms ($R^{14} + R^{15} + R^{16}$) is not greater than 18; or

X is $-CR^{17}=CR^{18}R^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue, the geometry of the enol

US Serial No. 10/597404
Page 16 of 25

double bond is E or Z, and the sum of all carbon atoms ($R^{17} + R^{18} + R^{19}$) is not greater than 18; or

X is $-CR^{17}=CR^{18}R^{19}$, wherein R^{17} , R^{18} and R^{19} are independently hydrogen or a C_1 - C_{18} hydrocarbon residue containing one or more atoms/groups selected from O, N and C(O), the geometry of the enol double bond is E or Z, and the sum of all carbon atoms ($R^{17} + R^{18} + R^{19}$) is not greater than 18;

R^{12} and R^{13} are independently hydrogen, C_1 - C_6 alkyl, C_1 - C_6 alkoxy residue, $-NO_2$, $-NH_2$, $-NHCO_2CH_3$, $-N(C_1-C_6 \text{ alkyl})_2$, $-N(\text{hydroxyalkyl})_2$, $-NHC(O)-(C_1-C_8 \text{ alkyl})$ or $-NHC(O)-(C_3-C_8 \text{ aryl})$; or

R^{12} and R^{13} are attached at the positions C(vi,vii), C(vii,viii), or C(viii,ix), and form together with the carbon atoms to which they are attached a dioxolane ring or a dioxane ring;

R^{11} in ii- or iii-position is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_3 - C_6 cycloalkyl, or $-CN$.

33.(withdrawn) A compound according to claim 32 wherein n is 1 and R^2 , R^3 and R^4 are hydrogen.

34.(withdrawn) A compound according to claim 32 wherein n is 1, R^2 and R^3 are hydrogen and R^4 is selected from methyl and $-CN$.

35.(withdrawn) A compound according to claim 32 wherein n is 1, R^2 and R^3 are hydrogen and R^4 is phenyl at C(3).

US Serial No. 10/597404
Page 17 of 25

36.(withdrawn) A compound according to claim 32 wherein n is 1, R² is hydrogen, R³ is at either C(6) to C(8) and R³ is selected from methyl, ethyl, propyl, isopropyl, methoxy, ethoxy and propyloxy, and R⁴ is selected from hydrogen, methyl, -CN, and phenyl at C(3).

37.(withdrawn) A compound according to claim 32 wherein n is 1, R² and R³ are methyl at position C(6,7), C(6,8), C(6,9), C(7,8), or C(8,9), and R⁴ is selected from hydrogen, -CN, and phenyl at C(3).

38.(withdrawn) A compound according to claim 32 wherein n is 1, R² and R³ are methoxy at C(7,9), and R⁴ is selected from hydrogen, -CN, and phenyl at C(3).

39.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen, methyl, -CN, and phenyl at C(3), R² is methyl at C(6) and R³ is isopropyl at C(9).

40.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen, methyl, -CN, and phenyl at C(3), R² is isopropyl at C(6) and R³ is methyl at C(9).

41.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, and R² and R³ are hydrogen.

42.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is hydrogen, and R³ is 7-methoxy.

43.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is hydrogen, and R³ is 6-methyl.

US Serial No. 10/597404
Page 18 of 25


44.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is hydrogen, and R³ is 7-methyl.

45.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is hydrogen, and R³ is 8-methyl.

46.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is hydrogen, and R³ is 6-tert-butyl.

47.(withdrawn) A compound according to claim 32 wherein n is 1, R⁴ is selected from hydrogen and methyl, R² is 6-tert-butyl, and R³ is 8-tert-butyl.

48.(withdrawn) A compound according to claim 32 wherein n is 1 and R is a monovalent residue of the formula (i)



wherein X, R¹¹, R¹² and R¹³ are defined according to claim 27.

49.(withdrawn) A compound according to claim 48 wherein R⁴=R¹¹, R³=R¹³ and R²=R¹².

50.(withdrawn) A process according to claim 31 wherein the compound of formula (I) hydrolysis in the presence of enzymes.